arXiv:1301.6300v1 [math.RT] 26 Jan 2013

LIFTING AUTOMORPHISMS OF QUOTIENTS OF ADJOINT REPRESENTATIONS

GERALD W. SCHWARZ

Abstract. Let V be the direct sum of r copies of the adjoint representation of the simple Lie group G. We assume that $r \geq 2$ and that $r \geq 3$ if G has rank 1. In [Sch12] we showed that any algebraic automorphism φ of the quotient $Z := V/\!\!/G$ lifts to a mapping $\Phi\colon V \to V$ which sends the fiber over $z \in Z$ to the fiber over $\varphi(z)$. (Most cases were already handled in Kuttler [Kut11].) Left open was the question of whether or not one can choose Φ to have some kind of equivariance property or whether one can choose Φ to be an automorphism of V. We show that any holomorphic automorphism φ of Z lifts to a holomorphic automorphism Φ of V such that $\Phi(gv) = \sigma(g)\Phi(v)$ for every $v \in V$ and $g \in G$ where σ is an automorphism of G. The key to the proof is showing that every algebraic differential operator on Z is induced by an invariant algebraic differential operator on V.

1. Introduction

Our base field is $\mathbb{C}$, the field of complex numbers. Let G be a complex reductive group and V a G-module. We denote the algebra of polynomial functions on V by $\mathcal{O}(V)$. For the following, we refer to [Kra84], [Lun73] and [VP89]. By Hilbert, the algebra $\mathcal{O}(V)^G$ is finitely generated, so that we have a quotient variety $Z := V/\!\!/G$ with coordinate ring $\mathcal{O}(Z) = \mathcal{O}(V)^G$. Let $\pi\colon V \to Z$ denote the morphism dual to the inclusion $\mathcal{O}(V)^G \subset \mathcal{O}(V)$. Then π sets up a bijection between the points of Z and the closed orbits in V. If Gv is a closed orbit, then the isotropy group $H = G_v$ is reductive. The *slice representation of H at v* is its action on N_v where N_v is an H-complement to $T_v(Gv)$ in $T_v(V) \simeq V$. Let $Z_{(H)}$ denote the points of Z such that the isotropy groups of the corresponding closed orbits are in the conjugacy class (H) of H. The $Z_{(H)}$ give a finite stratification of Z by locally closed smooth subvarieties. In particular, there is a unique open stratum $Z_{(H)}$, the *principal stratum*, which we also denote by Z_{pr}. We call H a *principal isotropy group* and any associated closed orbit a *principal orbit* of G.

As shorthand for saying that V has finite principal isotropy groups we say that V has FPIG. If V has FPIG, then there is an open set of closed orbits and a closed orbit is principal if and only if the slice representation of its isotropy group is trivial. Set $V_{\text{pr}} := \pi^{-1}(Z_{\text{pr}})$. We say that V is *k-principal* if V has FPIG and $\operatorname{codim} V \setminus V_{\text{pr}} \geq k$.

For X an affine variety let $\operatorname{Aut}(X)$ denote the automorphisms of X and let $\operatorname{Aut}_{\mathcal{H}}(X)$ denote the holomorphic automorphisms of X. Let $\varphi \in \operatorname{Aut}_{\mathcal{H}}(Z)$. We say that a holomorphic map $\Phi\colon V \to V$ is a *lift of φ* if $\pi \circ \Phi = \varphi \circ \pi$. Equivalently, Φ maps the fiber $\pi^{-1}(z)$ to the fiber $\pi^{-1}(\varphi(z))$, $z \in Z$. Let σ be an automorphism of G. We say that Φ is *σ-equivariant* if $\Phi(gv) = \sigma(g)\Phi(v)$ for all $v \in V$, $g \in G$.

Our main interest is in the following case. Let $V = \oplus_{i=1}^{d} r_i \mathfrak{g}_i$ where the $\mathfrak{g}_i$ are simple Lie algebras. Let G_i denote the adjoint group of $\mathfrak{g}_i$ and let G denote the product of the G_i. We assume that $r_i \geq 2$ for all i and that $r_i \geq 3$ if $\mathfrak{g}_i \simeq \mathfrak{sl}_2$. These conditions are necessary and sufficient for (V, G) to be 2-principal and for any $\varphi \in \operatorname{Aut}(Z)$ to preserve the stratification of Z given by the $Z_{(H)}$ [Sch13, Theorem 1.2, Proposition 3.1].

In [Sch12] we show that V has the following lifting property.

2010 *Mathematics Subject Classification.* 20G20, 22E46, 57S15.

Key words and phrases. differential operators, automorphisms, quotients.

Theorem 1.1. *Let V be as above. Then any $\varphi \in \operatorname{Aut}(Z)$ lifts to V.*

Most cases of the theorem were first established by Kuttler [Kut11]. This also holds for Corollary 1.3 below with φ algebraic. The lift given in the theorem is not necessarily an automorphism of V nor is it necessarily σ-equivariant for some σ.

The main result of this note is the following.

Theorem 1.2. *Let V be as above. Then for every $\varphi \in \operatorname{Aut}_{\mathcal{H}}(Z)$ there is an automorphism σ of G and a σ-equivariant $\Phi \in \operatorname{Aut}_{\mathcal{H}}(V)$ which lifts φ.*

Corollary 1.3. *Let $\varphi \in \operatorname{Aut}_{\mathcal{H}}(Z)$. Then there is an automorphism σ of G such that φ sends $Z_{(H)}$ to $Z_{(\sigma(H))}$ for every stratum $Z_{(H)}$ of Z.*

In §2 we recall two conditions sufficient for elements of $\operatorname{Aut}_{\mathcal{H}}(Z)$ to lift to σ-equivariant elements of $\operatorname{Aut}_{\mathcal{H}}(V)$. One of the conditions follows from Theorem 1.1 and the other is that one can lift algebraic differential operators on Z to invariant algebraic differential operators on V. In §3 we recall sufficient conditions for lifting differential operators and in §4 we show that V satisfies these conditions, giving a proof of Theorem 1.2.

2. Lifting automorphisms

Let G be reductive and V a G-module. We have a grading of $\mathcal{O}(Z) \simeq \mathcal{O}(V)^G$ induced from the grading of $\mathcal{O}(V)$, hence the scalar $\mathbb{C}^*$-action on V induces an action of $\mathbb{C}^*$ on Z. One sees that $t \cdot \pi(v) = \pi(t \cdot v)$ for $v \in V$ and $t \in \mathbb{C}^*$. Let $\operatorname{Aut}_{\mathrm{q}\ell}(Z)$ denote the elements of $\operatorname{Aut}(Z)$ whose action on $\mathcal{O}(Z)$ preserves the grading. Then $\operatorname{Aut}_{\mathrm{q}\ell}(Z)$ is a linear algebraic group. We say that V has the *lifting property* if every $\varphi \in \operatorname{Aut}_{\mathrm{q}\ell}(Z)$ has a lift $\Phi\colon V \to V$. If Φ lifts $\varphi \in \operatorname{Aut}_{\mathrm{q}\ell}(Z)$, then so does $\Phi'(0)$, so that liftable elements of $\operatorname{Aut}_{\mathrm{q}\ell}(Z)$ admit linear lifts.

Let $\mathcal{D}^k(V)$ (resp. $\mathcal{D}^k(Z)$) denote the algebraic differential operators on V (resp. Z) of order at most k (see [Sch95, §3]). Then we have a morphism $\pi_*\colon \mathcal{D}^k(V)^G \to \mathcal{D}^k(Z)$ where $\pi_*(P)(f) = P(\pi^*(f))$ for $P \in \mathcal{D}^k(V)^G$ and $f \in \mathcal{O}(Z) \simeq \mathcal{O}(V)^G$.

Definition 2.1. We say that the G-module V is *admissible* if

1. V is 2-principal.
2. $\pi_*\colon \mathcal{D}^k(V)^G \to \mathcal{D}^k(Z)$ is surjective for all k.

Strange as it may seem, admissibility has an application to the problem of lifting automorphisms of Z. From [Sch12, Theorems 1.3, 1.4] we have the following.

Theorem 2.2. *Suppose that V is admissible and has the lifting property. Then for every $\varphi \in \operatorname{Aut}_{\mathcal{H}}(Z)$ there is an automorphism σ of G and a σ-equivariant $\Phi \in \operatorname{Aut}_{\mathcal{H}}(V)$ which lifts φ.*

The proof of the theorem is briefly the following. Let $\varphi \in \operatorname{Aut}_{\mathcal{H}}(Z)$ and define $\varphi_t(z) = t^{-1} \cdot \varphi(t \cdot z)$ for $t \in \mathbb{C}^*$, $z \in Z$. Admissibility implies that φ_t extends to a holomorphic family on $\mathbb{C}$ with $\varphi_0 \in \operatorname{Aut}_{\mathrm{q}\ell}(Z)$. We can lift φ_0 to $\Phi_0 \in \operatorname{End}(V)$ by assumption. One shows that $\Phi_0 \in \mathrm{GL}(V)$ and that Φ_0 normalizes G, hence conjugation by Φ_0 gives an automorphism σ of G and Φ_0 is σ-equivariant. Then $\varphi_0^{-1}\varphi_t$ is an isotopy which is the identity of Z when $t = 0$. Thus $\varphi_0^{-1}\varphi_t$ is obtained by integrating a holomorphic vector field A_t on $\mathbb{C} \times Z$ and admissibility shows that we can lift A_t to an invariant holomorphic vector field B_t on $\mathbb{C} \times V$. Then integrating B_t from time 0 to time 1 gives an equivariant holomorphic lift Ψ of $\varphi_0^{-1}\varphi$ and $\Phi_0\Psi$ is a σ-equivariant automorphism lifting φ.

In the next section we recall sufficient conditions for admissibility.

3. Modularity

We need to recall some notions and results from [Sch94, Sch95]. Let H be a linear algebraic group and Y an H-variety. Let $Y_{(n)}$ or $(Y,H)_{(n)}$ denote the points of Y with isotropy group of dimension n. We define the *modularity of* Y, $\operatorname{mod}(Y,H)$, by

$$\operatorname{mod}(Y,H)=\sup_{n\geq 0}(\dim Y_{(n)}-\dim H+n)$$

where, as usual, the dimension of the empty set is $-\infty$. If $Y_{(n)}$ is irreducible, then $\dim Y_{(n)}-\dim H+n$ is the transcendence degree of $\mathbb{C}(Y_{(n)})^H$. We have that

$$\operatorname{mod}(Y,H)=\dim Y-\dim H+\sup_{n\geq 0}(n-\operatorname{codim} Y_{(n)}). \tag{3.0.1}$$

Now let X be an H-variety where $X_{(0)}\neq\emptyset$. We say that X is *k-modular* if $\operatorname{mod}(X\setminus X_{(0)},H)\leq\dim X-\dim H-k$. Equivalently, $\operatorname{codim} X_{(n)}\geq n+k$ for all $n\geq 1$. Let us assume from now on that H is reductive and that X is an affine H-variety with FPIG. Then $X_{(0)}\neq\emptyset$ and $\dim X-\dim H=\dim X/\!\!/H$ so that X is k-modular if and only if $\operatorname{mod}(X\setminus X_{(0)},H)\leq\dim X/\!\!/H-k$.

There are geometric interpretations of X being 0-modular. From now on assume that X is smooth. Let $A_1,\dots,A_r$ be a basis of the Lie algebra of H. We may consider the A_i as vector fields $\tilde{A}_i$ on X, hence as functions f_{A_i} on T^*X. Then X is 0-modular if and only if the f_{A_i} are a regular sequence in $\mathcal{O}(T^*X)$, i.e., their set of common zeroes has codimension r. Another interpretation is the following. Let $\mu\colon T^*X\to\mathfrak{h}^*$ be the moment mapping which sends $\xi\in T^*_xX$ to the element of $\mathfrak{h}^*$ whose value on $A\in\mathfrak{h}$ is $\tilde{A}_i(x)(\xi)$. Then X is 0-modular if and only if all the fibers of μ have codimension r [Sch95, Remark 8.6].

Remark 3.1. Note that if X is k-modular for any $k\geq 0$, then $\operatorname{mod}(X,H)=\dim X-\dim H$.

Suppose that X is 2-principal. Then we can find invariant functions h_1 and h_2 which vanish on $X\setminus X_{\mathrm{pr}}$ and are a regular sequence in $\mathcal{O}(X)$. If X is also 2-modular, then $h_1,h_2,f_{A_1},\dots,f_{A_r}$ is a regular sequence in $\mathcal{O}(T^*X)$ [Sch95, Lemma 9.7]. Then we can apply [Sch95, Proposition 8.15]:

Proposition 3.2. *Suppose that there are h_1, h_2 as above such that h_1, h_2 and the f_{A_i} are a regular sequence in $\mathcal{O}(T^*X)$. Then $\pi_*\colon\mathcal{D}^k(X)^H\to\mathcal{D}^k(X/\!\!/H)$ is surjective for every $k\geq 0$.*

Corollary 3.3. *Suppose that X is 2-principal and 2-modular. Then $\pi_*\colon\mathcal{D}^k(X)^H\to\mathcal{D}^k(X/\!\!/H)$ is surjective for every $k\geq 0$.*

In the next section we will show that our V of interest is 2-modular. We will need the following result. Recall that a group action is *almost effective* if the kernel of the action is finite.

Lemma 3.4. *Let H be connected reductive. Let T denote the connected center of H and let H_s denote $[H,H]$. Let $W:=W'\oplus W''$ be a direct sum of H-modules where T acts trivially on W'. Assume that the action of T on W'' is almost effective. If (W',H_s) is 0-modular, then $\operatorname{mod}(W,H)=\dim W-\dim H$.*

Proof. By Vinberg [Vin86], for every $q\geq 0$, $\operatorname{codim}(W'',T)_{(q)}\geq q$. Let $w=(w',w'')$ where $w'\in W'$ and $w''\in(W'',T)_{(q)}$. If $\dim H_w=p$, then $p\geq q$ and $\dim(H_s)_{w'}=p'-q$ for $p'\geq p$. Thus

$$\begin{aligned}
&\sup_{p\geq q}(p-\operatorname{codim}((W'\times(W'',T)_{(q)},H)_{(p)})\\
\leq&\sup_{p'\geq q}(p'-q-\operatorname{codim}((W',H_s)_{(p'-q)})+q-\operatorname{codim}(W'',T)_{(q)}\\
\leq&\sup_{p'\geq 0}(p'-\operatorname{codim}(W',H_s)_{(p')})=0.
\end{aligned}$$

Since $q \leq p$ is arbitrary, we see from (3.0.1) that $\operatorname{mod}(W, H) = \dim W - \dim H$. $\square$

4. Proof of the main theorem

We return to the study of the G-module $V = \oplus_{i=1}^d r_i\mathfrak{g}_i$. Recall that V is 2-principal and it has the lifting property by Theorem 1.1. Thus, by Corollary 3.3, we only need to show that V is 2-modular. It is easy to see that V is 2-modular if and only if each $r_i\mathfrak{g}_i$ is 2-modular, so we may reduce to the case that G is simple. The case $G = \mathrm{PSL}_2$ follows from [Sch95, Theorem 11.9], so we only have to consider the case where the rank of G is at least two. Moreover, we need only consider the case of two copies of $\mathfrak{g}$ since increasing the number of copies can only increase the modularity [Sch95, Proposition 11.5].

Let $v \in V \setminus V_{(0)}$. Then $\dim G_v > 0$ and there is an injective homomorphism $\lambda : H \to G_v$ where H is a copy of the multiplicative group $\mathbb{C}^*$ or the additive group $\mathbb{C}^+$. First consider the case where $H = \mathbb{C}^*$. Acting by an element of G, we may assume that $\operatorname{Im}\lambda \subseteq T$, where T denotes a maximal torus of G. Identify H with its image in T. There are only finitely many possible V^H. Let N_H denote $C_G(H)$. Then $\operatorname{mod}(G \cdot V^H, G)$ is bounded by $\operatorname{mod}(V^H, N_H)$.

Now suppose that $H = \mathbb{C}^+$. By the Jacobsen-Morozov theorem, λ extends to an injective homomorphism, also called λ, from SL_2 or PSL_2 to G. We identify H with its image in G and we let S denote $\lambda(\mathrm{SL}_2)$ or $\lambda(\mathrm{PSL}_2)$ as the case may be. Up to conjugation in G, there are only finitely many possible λ. Let N_H denote $C_G(S) \times T''$ where T'' is the maximal torus of S. Then again we have the estimate that $\operatorname{mod}(G \cdot V^H, G) \leq \operatorname{mod}(V^H, N_H)$.

Since every point in $V \setminus V_{(0)}$ lies in some V^H, and since $\dim V/\!\!/G = \dim G$, our discussion above gives the following.

Lemma 4.1. *Suppose that* $\operatorname{mod}(V^H, N_H) \leq \dim G - 2$ *for all H of dimension one. Then V is 2-modular.*

The next two propositions show that we always have $\operatorname{mod}(V^H, N_H) \leq \dim G - 2$, hence they complete our proof of Theorem 1.2.

Proposition 4.2. *Suppose that $H = \mathbb{C}^*$. Then* $\operatorname{mod}(V^H, N_H) \leq \dim G - 2$.

Proof. As above, we may assume that H is contained in a maximal torus T of G. The action of N_H on V^H is twice the adjoint representation of N_H. A maximal torus of N_H is T, and the connected center of N_H is a subtorus T' of T containing H. Then N_H/T' is semisimple, so its action on V^H is a trivial representation plus the sum of twice the adjoint representation of each simple component K_i of N_H/T'. By induction on rank, $2\mathfrak{k}_i$ is 2-modular if K_i has rank at least two. If $\mathfrak{k}_i \simeq \mathfrak{sl}_2$, then $2\mathfrak{k}_i$ is 1-modular. It then follows from Remark 3.1 that

$$\operatorname{mod}(V^H, N_H) = \dim V^H - \dim N_H + \dim T' = \dim \mathfrak{g}^H + \dim T'.$$

Let $\alpha_1, \dots, \alpha_k$ be the positive roots of $\mathfrak{g}$ which are nontrivial when restricted to T'. As T'-module,

$$\mathfrak{g} = \theta_m + \bigoplus_{i=1}^{k}(\mathfrak{g}_{\alpha_i} + \mathfrak{g}_{-\alpha_i})$$

where $m = \dim \mathfrak{g}^H$ and θ_m denotes the trivial module of dimension m. We need to show that $\dim \mathfrak{g} - 2 - (m + \dim T')$ is nonnegative, i.e., that $-2 - \dim T' + 2k \geq 0$. Since the action of T' on $\mathfrak{g}$ is effective, $k \geq \dim T'$. Thus, if $\dim T' \geq 2$, we obtain the desired estimate. The only problem is the case that $T' = H$ has dimension one. Suppose that only one positive root α of G acts nontrivially on H. We may assume that α is simple. Then, since G has rank at least two, there is a simple root β (acting trivially on H) such that $\alpha + \beta$ is a root. Then $\alpha + \beta$ acts nontrivially on H and we have a contradiction. Hence $k \geq 2$ and $\operatorname{mod}(V^H, N_H) \leq \dim G - 2$. $\square$

Proposition 4.3. *Suppose that* $H = \mathbb{C}^+$. *Then* $\operatorname{mod}(V^H, N_H) \leq \dim V/\!\!/G - 2$.

Proof. Let $H \subset S \subset G$ be as discussed above. As S-module, $\mathfrak{g} = \sum_i m_i R_i$ where R_i denotes the space of binary forms of degree i and $m_i \geq 0$ is the multiplicity of R_i. The fixed points of H in each R_i have dimension one, so that V^H has dimension $2\sum_i m_i$. The Lie algebra of N_H has dimension $m_0 + 1$ and the action of $C_G(S)$ on $2\mathfrak{g}^S$ is twice its adjoint representation. Let T' be the connected center of the identity component of $C_G(S)$. Recall that T'' is the maximal torus of S. Now T' acts on $m_i R_i$ via a homomorphism to GL_{m_i}, hence T' acts almost effectively on $\oplus_{i\geq 1} m_i R_i^H$. Since T' commutes with S, it acts trivially on $\mathfrak{s} \subset m_2 R_2$. However, T'' acts nontrivially on $\mathfrak{s}^H$. Thus $T' \times T''$ acts almost effectively on V^H. Using induction on rank and Lemma 3.4, we get that $\operatorname{mod}(V^H, N_H)$ is $\dim V^H - \dim C_G(S) - 1$. Now $\dim V^H$ is $2(m_0 + \sum_{i\geq 1} m_i)$ and $\dim C_G(S) = m_0$. We need that $\dim V^H - m_0 - 1 \leq \dim \mathfrak{g} - 2$. Hence we need

$$2(m_0 + \sum_{i\geq 1} m_i) - m_0 - 1 \leq \sum_{i\geq 0} (i+1) m_i - 2 \text{ ,i.e., } 0 \leq -1 + \sum_{i\geq 1} (i-1) m_i.$$

Since $\mathfrak{s} \simeq R_2 \subset \mathfrak{g}$, we have that $m_2 \geq 1$, giving us the desired inequality. ☐

References

[Kra84] Hanspeter Kraft, *Geometrische Methoden in der Invariantentheorie*, Aspects of Mathematics, D1, Friedr. Vieweg & Sohn, Braunschweig, 1984.

[Kut11] J. Kuttler, *Lifting automorphisms of generalized adjoint quotients*, Transformation Groups **16** (2011), 1115–1135.

[Lun73] Domingo Luna, *Slices étales*, Sur les groupes algébriques, Soc. Math. France, Paris, 1973, pp. 81–105. Bull. Soc. Math. France, Paris, Mémoire 33.

[Sch94] Gerald W. Schwarz, *Differential operators on quotients of simple groups*, J. Algebra **169** (1994), no. 1, 248–273.

[Sch95] ______, *Lifting differential operators from orbit spaces*, Ann. Sci. École Norm. Sup. (4) **28** (1995), no. 3, 253–305.

[Sch12] ______, *Quotients, automorphisms and differential operators*, http://arxiv.org/abs/1201.6369 (2012).

[Sch13] ______, *Vector fields and Luna strata*, J. Pure and Applied Algebra **217** (2013), 54–58.

[Vin86] È. B. Vinberg, *Complexity of actions of reductive groups*, Funktsional. Anal. i Prilozhen. **20** (1986), no. 1, 1–13, 96.

[VP89] È. B. Vinberg and V. L. Popov, *Invariant theory*, Algebraic geometry, 4 (Russian), Itogi Nauki i Tekhniki, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989, pp. 137–314, 315.

Department of Mathematics, Brandeis University, Waltham, MA 02454-9110
E-mail address: `schwarz@brandeis.edu`